

MT
1785

Master's Thesis

Diversity and Endemism of Soybean Rhizobia between Southern Japan and
Central Luzon, Philippines

Academic year: 2015

Graduate School of Agriculture,
University of Miyazaki

Course: International Course of Agriculture

Name: Maria Luisa T. Mason

Principal advisor

Name: Yuichi Saeki 印

Date of acceptance

01 / 29 / 2016
Day/Month/Year

Graduate School of Agriculture

Table of Contents

	Page
List of Tables	iv
List of Figures	v
Chapter 1 General introduction.....	1
Chapter 2 Diversity of soybean rhizobia through PCR-RFLP analysis of the 16S-23S rRNA gene internal transcribed spacer (ITS) region and <i>rpoB</i> housekeeping gene	
2.1 Introduction.....	8
2.2 Materials and Methods.....	10
2.2.1 Soil analysis (pH and EC).....	10
2.2.2 Random sampling of representative samples.....	11
2.2.3 Isolation of soybean rhizobia.....	11
2.2.4 Incubation.....	12
2.2.5 DNA extraction.....	13
2.2.6 PCR amplification of 16S-23S rRNA gene ITS region.....	13
2.2.7 PCR amplification of <i>rpoB</i> housekeeping gene.....	14
2.2.8 Confirmation of amplified PCR product.....	16
2.2.9 Restriction enzyme treatment of PCR amplified products for ITS region (RFLP analysis).....	16

2.2.10	Restriction enzyme treatment of PCR amplified products for <i>rpoB</i> housekeeping gene (RFLP analysis).....	16
2.2.11	Electrophoresis of the solution digested by restriction enzymes.....	17
2.2.12	Construction of phylogenetic tree.....	17
2.3	Results.....	20
2.3.1	Morphological growth of isolates.....	21
2.3.2	16S-23S rRNA gene ITS region.....	23
2.3.3	<i>rpoB</i> housekeeping gene.....	23
2.4	Discussion.....	28
Chapter 3	Sequence analysis of the 16S rRNA gene, 16S-23S rRNA gene ITS region, and <i>rpoB</i> housekeeping gene between Southern Japan and Central Luzon, Philippines	
3.1	Introduction.....	32
3.2	Materials and Methods.....	34
3.2.1	Sample selection.....	34
3.2.2	PCR amplification of isolates and reference strains....	34
3.2.3	Purification of PCR amplified products.....	35
3.2.4	Sample preparation for sequence analysis.....	35
3.2.5	Sequence alignment and phylogenetic tree construction.....	36
3.2.6	Multidimensional scaling analysis.....	37

3.3	Results.....	37
3.3.1	16S rRNA gene sequence analysis.....	37
3.3.2	16S-23S rRNA gene ITS region sequence analysis....	39
3.3.3	<i>rpoB</i> housekeeping gene sequence analysis.....	39
3.3.4	Multidimensional scaling analysis.....	42
3.4	Discussion.....	46
Chapter 4	General discussion	51
References	59
Acknowledgment	74
Supplementary data	75

List of Tables

Table 1	Some information on the study sites.....	20
Table 2	Summary of RFLP analysis in Kumamoto, Okinawa, and C. Luzon, Philippines.....	20
Table 3	Morphological characteristics of the colonies observed for the 18 isolates.....	22
Table 4	List of sequence primers used in this study.....	36
Table 5	Distance matrix of 16S-23S rRNA gene ITS region from the sequence analysis of 18 isolates and <i>B.elkanii</i> USDA reference strains.....	42
Table 6	Distance matrix of <i>rpoB</i> housekeeping gene from the sequence analysis of 18 isolates and <i>B.elkanii</i> USDA reference strains.....	42
Table 7	Comparison between RFLP and sequence analysis of the 18 isolates.....	50

List of Figures

Figure 1	Flowchart of activities conducted.....	19
Figure 2	Restriction enzymes treatment band patterns of ITS region for the 18 isolates and 11 USDA <i>Bradyrhizobium</i> reference strains.....	25
Figure 3	Restriction enzymes treatment band patterns of <i>rpoB</i> housekeeping gene for the 18 isolates.....	26
Figure 4	Restriction enzymes treatment band patterns of <i>rpoB</i> housekeeping gene of <i>Bradyrhizobium</i> USDA reference strains.....	27
Figure 5	Average monthly temperature of the three locations.....	30
Figure 6	Complete genome sequence of <i>Bradyrhizobium japonicum</i> USDA110.....	31
Figure 7	Phylogenetic tree based on sequence analysis of 16S rRNA gene.....	38

Figure 8	Phylogenetic tree based on sequence analysis of 16S-23S rRNA gene ITS region.....	40
Figure 9	Phylogenetic tree based on sequence analysis of <i>rpoB</i> housekeeping gene.....	41
Figure 10	Multidimensional scaling (MDS) plot of 16S-23S rRNA gene ITS region showing similarities in the nucleotide sequence between <i>B. elkanii</i> USDA reference strains and the 18 isolates.....	43
Figure 11	Multidimensional scaling (MDS) plot of 16S-23S rRNA gene ITS region showing similarities in the nucleotide sequence between <i>B. elkanii</i> USDA reference strains (excluding USDA Be94) and the 18 isolates.....	44
Figure 12	Multidimensional scaling (MDS) plot of <i>rpoB</i> housekeeping gene showing similarities in the nucleotide sequence between <i>B. elkanii</i> USDA reference strains and the 18 isolates.....	45
Figure 13	Multidimensional scaling (MDS) plot of <i>rpoB</i> housekeeping gene showing similarities in the nucleotide sequence between <i>B. elkanii</i> USDA reference strains (excluding Be94) and the 18 isolates.....	45

REFERENCES

- Ade'kambi, T., Shinnick, T., Raoult, D., & Drancourt, M. (2008). Complete *rpoB* gene sequencing as a suitable supplement to DNA–DNA hybridization for bacterial species and genus delineation. *Int. J. Syst. Evol. Microbiol.*, 58, 1807–1814. doi: 10.1099/ijs.0.65440-0
- Adhikari, D., Kaneto, M., Itoh, K., Suyama, K., Pokharel, B.B., & Gaihre, Y.K. (2012). Genetic diversity of soybean-nodulating rhizobia in Nepal in relation to climate and soil properties. *Plant Soil*, 357, 131–145. doi: 10.1007/s11104-012-1134-6
- Ahmad, S., Mokaddas, E., & Fares, E. (2002). Characterization of *rpoB* mutations in rifampin-resistant clinical *Mycobacterium tuberculosis* isolates from Kuwait and Dubai. *Diagn. Microbiol. Infect. Dis.*, 44, 245–252.
- Alam, F., Bhuiyan, M.A., Alam, S.S., Waghmode, T.R., Kim, P.J., & Lee, Y.B. (2015). Effect of *Rhizobium* sp. BARIRGm901 inoculation on nodulation, nitrogen fixation and yield of soybean (*Glycine max*) genotypes in gray terrace soil. *Biosci. Biotechnol. Biochem.*, 79 (10), 1660-8. doi: 10.1080/09168451.2015.1044931
- Ansari, P.G., Rao, D.L.N., & Pal, K.K. (2013). Diversity and phylogeny of soybean rhizobia in central India. *Ann. Microbiol.*, 64, 1553–1565. doi: 10.1007/s13213-013-0799-2
- Asad, S., Malik, K.A., & Hafeez, F.Y. (1991). Competition between inoculated and indigenous *Rhizobium/Bradyrhizobium* spp. strains for nodulation of grain and fodder legumes in Pakistan. *Biol. Fert. Soils*, 12 (2), 107-111.

- Aserse, A.A., Räsänen, L.A., Assefa, F., Hailemariam, A., & Lindström, K. (2012). Phylogeny and genetic diversity of native rhizobia nodulating common bean (*Phaseolus vulgaris* L.) in Ethiopia. *Syst. Appl. Microbiol.*, 35, 120–131. doi: 10.1016/j.syapm.2011.11.005
- Baraúna, A.C., de Silva, K., Pereira, G.M.D., Kaminski, P.E., Perin, L., & Zilli, J.E. (2014). Diversity and nitrogen fixation efficiency of rhizobia isolated from nodules of *Centrolobium paraense*. *Pesq. agropec. bras., Brasília*, 49 (4), 296-305. doi: 10.1590/S0100-204X2014 000400008
- Berrada, H. & Fikri-Benbrahim, K. (2014). Taxonomy of the Rhizobia: Current Perspectives. *Br. Microbiol. Res. J.*, 4(6), 616-639.
- Bureau of Agricultural Statistics (BAS). Retrieved from <http://countrystat.psa.gov.ph/>
- Cheng, Q. (2008). Perspectives in biological nitrogen fixation research. *J. Integr. Plant Biol.*, 50 (7), 784–796. doi: 10.1111/j.1744-7909.2008.00700.x
- Chriki-Adeeb, R. & Chriki, A. (2015). Bayesian Phylogenetic Analysis of Rhizobia Isolated From Root-Nodules of Three Tunisian Wild Legume Species of the Genus *Sulla*. *J. Phylogen. Evolution. Biol.*, 3 (2). <http://dx.doi.org/10.4172/2329-9002.1000149>
- Cole, M.A. & Elkan, G.H. (1973) Transmissible resistance to penicillin G, neomycin, and chloramphenicol in *Rhizobium japonicum*. *Antimicrob. Ag. Chemother.*, 4, 248-253.
- Degefu, T., Wolde-meskel, E., & Frostegård, A. (2013). Phylogenetic diversity of *Rhizobium* strains nodulating diverse legume species growing in Ethiopia. *Syst. Appl. Microbiol.*, 36, 272–280. doi.org/10.1016 /j.syapm.2013.03.004

- Delamuta, J.R.M., Ribeiro, R.A., Menna, P., Bangel, E.V., & Hungria, M. (2012). Multilocus Sequence Analysis (MLSA) Of *Bradyrhizobium* Strains: Revealing High Diversity of Tropical Diazotrophic Symbiotic Bacteria. *Braz. J. Microbiol.*, 698-710.
- Devine, T.E. & Kuykendall, L.D. (1996). Host genetic control of symbiosis in soybean (*Glycine max* L.). *Plant Soil*, 186, 173 –187.
- Dugje, I.Y., Omoigui, L.O., Ekeleme, F., Bandyopadhyay, R., Lava Kumar, P., & Kamara, A.Y. (2009). Farmers' Guide to Soybean Production in Northern Nigeria. *International Institute of Tropical Agriculture*, Ibadan, Nigeria. 21 pp.
- Dwevedi, A. & Kayastha, A. (2011). Soybean: a Multifaceted Legume with Enormous Economic Capabilities, Soybean - Biochemistry, Chemistry and Physiology, Prof. Tzi-Bun Ng (Ed.), ISBN: 978-953-307- 219-7, *InTech*.
- Food and Agriculture Organization (FAO) of the United Nations. FAOSTAT. www.faostat.org.
- Fuhrmann, J. (1990). Symbiotic effectiveness of indigenous soybean bradyrhizobia as related to serological, morphological, rhizobitoxine, and hydrogenase phenotypes. *Appl. Environ. Microbiol.*, (56), 224–229.
- Gamo, T., Itoh, A., Sawazaki, A., Manguiat, I., & Mendoza, D. (1990). Rhizobia collected from leguminous plants in the Philippines. *Bull. Natl. Inst. Agrobiol.*, Resour. 6: 111 – 129.
- Germano, M.G., Menna, P., Mostasso, F.L., & Hungria, M. (2006). RFLP analysis of the rRNA operon of a Brazilian collection of bradyrhizobial strains from 33 legume species. *Int. J. Syst. Evol. Microbiol.*, 56, 217–229. doi: 10.1099/ijs.0.02917-0

- Guimarães, A.A., Florentino, L.A., Almeida, K.A., Lebbe, L., Silva, K.B., Willems, A., Moreira, & F.M. (2015). High diversity of *Bradyrhizobium* strains isolated from several legume species and land uses in Brazilian tropical ecosystems. *Syst. Appl. Microbiol.*, 38, 433–441. doi.org/10.1016/j.syapm. 2015.06.006
- Hayashi, M., Saeki, Y., Haga, M., Harada, K., Kouchi, H., & Umehara, Y. (2012). *Rj* (*rj*) genes involved in nitrogen-fixing root nodule formation in soybean. *Breed Sci.*, 61(5), 544–553. doi:10.1270/jsbbs.61.544
- Hiraishi, A., Kamagata, Y. & Nakamura, K. (1995). Polymerase chain reaction amplification and restriction fragment length polymorphism analysis of 16S rRNA genes from Methanogens. *J. Ferment. Bioengineer.*, 79, 523 – 529.
- Htwe, A.Z., Yamakawa, T., Sarr, P.S., & Sakata, T. (2015). Diversity and distribution of soybean-nodulating bradyrhizobia isolated from major soybean-growing regions in Myanmar. *Afr. J. Microbiol. Res.*, 9 (43), 2183-2196. doi:10.5897/AJMR2015.7702
- Ikeda, S., Rallos, L.E.E., Okubo, T., Eda, S., Inaba, S., Mitsui, H. & Minamisawa, K. (2008). Microbial community analysis of field-grown soybeans with different nodulation phenotypes. *Appl. Environ. Microbiol.*, 74, 5704–5709. doi:10.1128/AEM.00833-08
- Ikeda, S., Okubo, T., Kaneko, T., Inaba, S., Maekawa, T., Eda, S., Sato, S., Tabata, S., Mitsui, H., & Minamisawa, K. (2010). Community shifts of soybean stem-associated bacteria responding to different nodulation phenotypes and N levels. *ISME J.*, 4, 315–326. doi:10.1128/AEM.00833-08
- Jordan, DC. 1982. Transfer of *Rhizobium japonicum* Buchanan 1980 to *Bradyrhizobium* gen. nov., a genus of slow-growing, root nodule bacteria from leguminous plants. *Int. J. Syst. Bacteriol.*, 32,136–139.

- Keyser, H. & Li, F. (1992). Potential for increasing biological nitrogen fixation in soybean. *Plant Soil*, 141, 119-135.
- Khamis, A., Colson, P., Raoult, D., & La Scola, B. (2003). Usefulness of *rpoB* Gene Sequencing for Identification of *Azotobacter* and *Besseyia* Species, Including a Strategy for Choosing Discriminative Partial Sequences. *Appl. Environ. Microbiol.*, 69 (11), 6740–6749. doi: 10.1128/AEM.69.11.6740–6749.2003
- Kimura, M. (1980). A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.*, 16, 111–120.
- Kober, M. dV., Saccol de Sa, E. L., Freire, J.R.J., & Giongo, A. (2004). Characterization of variants *Bradyrhizobium elkanii* and *B. japonicum* and symbiotic behaviour in soybeans. *Ciencia Rural, Santa Maria*, 34 (5), 1459-1464.
- Kuykendall, L.D., Saxena, B., Cevine, T.E., & Udell, S.E. (1992). Genetic diversity in *Bradyrhizobium* Jordan 1982 and a proposal for *Bradyrhizobium elkanii* sp. nov. *Can. J. Microbiol.*, 38, 501–505.
- Kvien, C.S., Ham, G.E., & Lambert, J.W. (1981). Recovery of Introduced *Rhizobium japonicum* Strains by Soybean Genotypes. *Agron. J.*, 73(5), 900-905. doi:10.2134/agronj1981.00021962007300050034x
- Lamphey, S., Ahiabor, B.D.K., Yeboah, S., & Asamoah, C. (2014). Response of soybean (*Glycine max*) to rhizobial inoculation and phosphorus application. *J. Exp. Biol. Agricul. Sci.*, 2 (1), 73-77. ISSN No. 2320 – 8694
- Li, Q.Q., Wang, E.T., Zhang, Y.Z., Zhang, Y.M., Tian, C.F., Sui, X.H., Chen, W.F., & Chen, W.X. (2011). Diversity and Biogeography of Rhizobia Isolated from

- Root Nodules of *Glycine max* Grown in Hebei Province, China. *Microbiol. Ecol.*, 61 (4), 917-931. doi:10.1007/s00248-011-9820-0
- Lin, D.X., Man, C.X., Wang, E.T., & Chen, W.X. (2007). Diverse rhizobia that nodulate two species of *Kummerowia* in China. *Arch. Microbiol.*, 188 (5), 495-50. doi: 10.1007/s00203-007-0271-4
- Loureiro, M., Kaschuk, G., Alberton, O., & Hungria, M. (2007). Soybean [*Glycine max* (L.) Merrill] rhizobial diversity in Brazilian oxisols under various soil, cropping, and inoculation managements. *Biol. Fertil. Soils*, 43, 665-674. doi: 10.1007/s00374-006-0146-x
- Man, C.X., Wang, H., Chen, W.F., Sun, X.H., Wang, E.T., & Chen, W.X. (2008). Diverse rhizobia associated with soybean grown in the subtropical and tropical regions of China. *Plant Soil*, 310 (1), 77-87. doi: 10.1007/s11104-008-9631-3
- Martens, M., Dawyndt, P., Coopman, R., Gillis, M., De Vos, P., & Willems, A. (2008). Advantages of multilocus sequence analysis for taxonomic studies: a case study using 10 housekeeping genes in the genus *Bradyrhizobium* (including former *Sinorhizobium*). *Int. J. Syst. Evol. Microbiol.*, 58, 200-214. doi:10.1099/ijs.0.653920
- Masuda, T. & Goldsmith, P. (2009). World Soybean Production: Area Harvested, Yield, and Long-Term Projections. *Int. Food Agribus. Manag. Rev.*, 12 (4), 143-162.
- McCammon, M.T., Gillette, J.S., Thomas, D.P., Ramaswamy, S.V., Graviss, E.A., Kreiswirth, B.N., Vijg, J., & Quiñones, T.N. (2005). Detection of *rpoB* mutations associated with rifampin resistance in *Mycobacterium tuberculosis* using denaturing gradient gel electrophoresis. *Antimicrob. Agents Chemother.*, 49(6), 2200-9. doi: 10.1128/AAC.49.6.2200-2209.2005

- Miller, R.H. & May, S. (1991). The Rhizosphere and Plant Growth: Legume inoculation: Success and failures. *Beltsville Symposia in Agr. Res.*, 14, 123-124. doi:10.1007/978-94-011-3336-4_27
- Minami, M., Yamakawa, T., Yamamoto, A., Akao, S., & Saeki, Y. (2009). Estimation of nodulation tendency among *Rj*-genotype soybeans using the bradyrhizobial community isolated from an Andosol. *Soil Sci. Plant Nutr.*, 55, 65–72. doi: 10.1111/j.1747-0765.2008.00333.x
- Minamisawa, K., Onodera, S., Tanimura, Y., Kobayashi, N., Yuhashi, K., & Kubota, M. (1997). Preferential nodulation of *Glycine max*, *Glycine soja* and *Macroptilium atropurpureum* by two *Bradyrhizobium* species *japonicum* and *elkanii*. *FEMS Microbiol. Ecol.*, 24, 49-56.
- Nelson, M. & Sadowsky, M. (2015). Secretion systems and signal exchange between nitrogen-fixing rhizobia and legumes. *Front. Plant Sci.*, 6: 491. doi: 10.3389/fpls.2015.00491
- Njira, K.O.W., Nalivata, P.C., Kanyama-Phiri, G.Y., & Lowole, M.W. (2013). An assessment for the need of soybean inoculation with *Bradyrhizobium japonicum* in some sites of Kasungu district, Central Malawi. *Int. J. Curr. Microbiol. App. Sci.*, 2 (8), 60-72. ISSN: 2319-7706
- Nguyen, M.T., Akiyoshi, K., Nakatsukasa, M., Saeki, Y., & Yokoyama, K. (2010). Multiple occupancy of nodules by nodulating rhizobia on field-grown soybeans with attendance of *Sinorhizobium* spp. *Soil Sci. Plant Nutr.*, 56 (3), 382 – 389. doi:10.1111/j.1747-0765.2010.00476.x
- Nzoué, A., Miche, L., Klonowska, A., Laguerre, G., de Lajudie, P., & Moulin, L. (2009). Multilocus sequence analysis of bradyrhizobia isolated from *Aeschynomene* species in Senegal. *Syst. Appl. Microbiol.*, 32, 400–412.

doi:10.1016/j.syapm.2009.06.002

- Patra, R.K., Pant, L.M., & Pradhan, K. (2012). Response of Soybean to Inoculation with Rhizobial Strains: Effect on Growth, Yield, N Uptake and Soil N Status. *World J. Agric. Sci.*, 8 (1), 51-54. ISSN 1817-3047
- Potts, J., Lynch, M., Wilkings, A., Huppé, G., Cunningham, M., & Voora, V. (2014). The State of Sustainability Initiatives Review 2014. Standards and the Green Economy. *International Institute for Sustainable Development (IISD) and the International Institute for Environment and Development (IIED)*. ISBN: 978-1-894784-45-0. 12, 253-273.
- Qian, J., Kwon, S.W., & Parker, M.A. (2003). rRNA and *nifD* phylogeny of *Bradyrhizobium* from sites across the Pacific Basin. *FEMS Microbiol. Lett.*, 219, 159-165. doi :10.1016/S0378-1097(03)00043-0
- Rajendhran, J. & Gunasekaran, P. (2011). Microbial phylogeny and diversity: Small subunit ribosomal RNA sequence analysis and beyond. *Microbiol. Res.*, 166, 99 – 110. doi: 10.1016/j.micres.2010.02.003
- Ribeiro, R.A., Barcellos, F.G., Thompson, F.L., & Hungria, M. (2009). Multilocus sequence analysis of Brazilian *Rhizobium* microsymbionts of common bean (*Phaseolus vulgaris* L.) reveals unexpected taxonomic diversity. *Res. Microbiol.*, 160, 297-306. doi:10.1016/j.resmic.2009.03.009
- Risal, C.P., Yokoyama, T., Ohkama-Ohtsu, N., Djedidi, S., & Sekimoto, H. (2010). Genetic diversity of native soybean bradyrhizobia from different topographical regions along the southern slopes of the Himalayan Mountains in Nepal. *Syst. Appl. Microbiol.*, 33, 416–425. doi:10.1016/j.syapm.2010. 06.008
- Rivas, R., Martens, M., de Lajudie, P., & Willems, A. (2009). Multilocus sequence analysis of the genus *Bradyrhizobium*. *Syst. Appl. Microbiol.*, 32, 101–110.

doi:10.1016/j.syapm.2008.12.005

- Ruiz Diaz, D.A., Pedersen, P., & Sawyer, J.E. (2009). Soybean Response to Inoculation and Nitrogen Application Following Long-Term Grass Pasture. *Crop Sci.*, 49 (3), 1058-1062. doi: 10.2135/cropsci2008.08.0510
- RStudio Team (2015). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA URL <http://www.rstudio.com/>.
- Sameshima, R., Isawa, T., Sadowsky, M.J., Hamada, T., Kasai, H., Shutsrirung, A., Mitsui, H. & Minamisawa, K. (2003). Phylogeny and distribution of extra-slow-growing *Bradyrhizobium japonicum* harboring high copy numbers of RS α , RS β and IS1631. *FEMS Microbiol. Ecol.*, 44, 191–202. doi:10.1016/S0168-6496(03)00009-6
- Sancholi, G., Mobasser, H.R., & Fanaei, H.R. (2015). Effect Inoculation of Soybean Cultivars with bacteria *Rhizobium japonicum* in Sistan. *Biological Forum – An Int. J.*, 7 (1), 552-558. ISSN No. 0975-1130
- Sanz-Sáez, A., Heath, K., Burke, P., & Ainsworth, E. (2015). Inoculation with an enhanced N₂O-fixing *Bradyrhizobium japonicum* strain (USDA110) does not alter soybean (*Glycine max* Merr.) response to elevated [CO₂]. *Plant Cell Environ.*, 38, 2589-2602. doi: 10.1111/pce.12577
- Saeki, Y., Aimi, N., Hashimoto, M., Tsukamoto, S., Kaneko, A., Yoshida, N., Nagatomo, Y., & Akao, S. (2004). Grouping of *Bradyrhizobium* USDA Strains by Sequence Analysis of 16S rDNA and 16S-23S rDNA Internal Transcribed Spacer Region. *Soil Sci. Plant Nutr.*, 50 (4), 517-525.
- Saeki, Y., Kaneko, A., Hara, T., Suzuki, K., Yamakawa, T., Nguyen, M.T., Nagatomo, Y., & Akao, S. (2005). Phylogenetic Analysis of Soybean-Nodulating Rhizobia Isolated from Alkaline Soils in Vietnam. *Soil Sci. Plant*

Nutr., 51 (7), 1043-1052.

- Saeki, Y., Aimi, N., Tsukamoto, S., Yamakawa, T., Nagatomo, Y., & Akao, S. (2006). Diversity and geographical distribution of indigenous soybean-nodulating bradyrhizobia in Japan. *Soil Sci. Plant Nutr.*, 52, 418–426. doi:10.1111/j.1747-0765.2006.00050.x
- Saeki, Y., Minami, M., Yamamoto, A., & Akao, S. (2008). Estimation of the bacterial community diversity of soybean-nodulating bradyrhizobia isolated from *Rj*-genotype soybeans. *Soil Sci. Plant Nutr.*, 54, 718–724. doi: 10.1111/j.1747-0765.2008.00300.x
- Saeki, Y., Ozumi, S., Yamamoto, A., Umehara, Y., Hayashi, M., & Sigua, G.C. (2010). Changes in population occupancy of bradyrhizobia under different temperature regimes. *Microbes Environ.*, 25, 309-312. doi:10.1264/jsme2.ME10128
- Saeki, Y. (2011). Characterization of soybean-nodulating rhizobial communities and diversity, p. 163–184. In A. Sudaric (ed.), Soybean- Molecular Aspects of Breeding. *InTech*, Croatia.
- Saeki, Y., Shiro, S., Tajima, T., Yamamoto, A., Sameshima-Saito, R., Sato, T., & Yamakawa, T. (2013). Mathematical Ecology Analysis of Geographical Distribution of Soybean-Nodulating Bradyrhizobia in Japan. *Microbes Environ.*, 28 (4), 470–478. doi:10.1264/jsme2.ME13079
- Saeki, Y. & Shiro, S. (2014). Comparison of Soybean-Nodulating Bradyrhizobia Community Structures Along North Latitude Between Japan and USA, Advances in Biology and Ecology of Nitrogen Fixation, Prof. Takuji Ohyama (Ed.), ISBN: 978-953-51-1216-7, *InTech*, doi:10.5772/57165

- Scholla, M.H. & Elkan, H.G. (1984). *Rhizobium fredii* sp.nov. a fast-growing species that effectively nodulates soybeans. *Int. J. Syst. Bacteriol.*, 34, 484–486.
- Sharma, M., Srivastava, K., & Sharma, S. (2010). Biochemical characterization and metabolic diversity of soybean rhizobia isolated from Malwa region of Central India. *Plant Soil Environ.*, 56 (8), 375–383.
- Shiro S, Yamamoto A, Umehara Y, Hayashi M, Yoshida N, Nishiwaki A, Yamakawa T, & Saeki Y. 2012. Effect of *Rj* genotype and cultivation temperature on the community structure of soybean-nodulating bradyrhizobia. *Appl. Environ. Microbiol.*, 78, 1243–1250. doi: 10.1128/AEM.06239-11
- Shiro, S., Matsuura, S., Saiki, R., Sigua, G.C., Yamamoto, A., Umehara, Y., Hayashi, M., & Saeki, Y. (2013). Genetic diversity and geographical distribution of indigenous soybean-nodulating bradyrhizobia in the United States. *Appl. Environ. Microbiol.*, 79, 3610– 3618. doi: 10.1128/AEM.00236-13
- Simon, Z., Mtei, K., Gessesse, A., & Ndakidemi, P. A. (2014). Isolation and Characterization of Nitrogen Fixing Rhizobia from Cultivated and Uncultivated Soils of Northern Tanzania. *Am. J. Plant Sci.*, 5, 4050-4067. <http://dx.doi.org/10.4236/ajps.2014.526423>
- Soe, K.M., & Yamakawa, T. (2013). Low-Density Co-Inoculation of Myanmar *Bradyrhizobium yuanmingense* MAS34 and *Streptomyces griseoflavus* P4 to Enhance Symbiosis and Seed Yield in Soybean Varieties. *Am. J. Plant Sci.*, 4, 1879-1892. <http://dx.doi.org/10.4236/ajps.2013.49231>
- Stackebrandt, E., & Goebel, B.M. (1994). Taxonomic note: a place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int. J. Syst. Bacteriol.*, 44, 846-849.

- Sulieman, S., & Tran, L.-S. P. (2014). Symbiotic Nitrogen Fixation in Legume Nodules: Metabolism and Regulatory Mechanisms. *Int. J. Mol. Sci.*, 15 (11), 19389–19393. <http://doi.org/10.3390/ijms151119389>
- Suresh, N., Singh, U.B., Arora, J., Pant, H., Seth, P., Sola, C., Rastogi, N., Samantaray, J.C., & Pande, J.N. (2006). *rpoB* gene sequencing and spoligotyping of multidrug-resistant *Mycobacterium tuberculosis* isolates from India. *Infect. Genet. Evol.*, 6, 474–483. doi:10.1016/j.meegid.2006.03.001
- Suzuki, K., Oguro, H., Yamakawa, T., Yamamoto, A., Akao, S., & Saeki, Y. (2008). Diversity and distribution of indigenous soybean - nodulating rhizobia in the Okinawa islands, Japan. *Soil Sci. Plant Nutr.*, 54, 237–246. doi: 10.1111/j.1747-0765.2007.00236.x
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., & Kumar, S. (2013). MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Mol. Biol. Evol.*, 30 (12), 2725–2729. doi:10.1093/molbev/mst197
- Tang, J., Bromfield, E.S.P., Rodrigue, N., Cloutier, S., & Tambong, J.T. (2012). Microevolution of symbiotic *Bradyrhizobium* populations associated with soybeans in east North America. *Ecol. Evol.*, 2, 2943–2961. doi: 10.1002/ece3.404
- Thies J.E., Bohlool, B.B., & Singleton, P.W. (1992). Environmental effects on competition for nodule occupancy between introduced and indigenous rhizobia and among introduced strains. *Can. J. Microbiol.*, 38, 493-500.
- Tringe, S. & Hugenholtz, P. (2008). A renaissance for the pioneering 16S rRNA gene. *Curr. Opin. Microbiol. – Genomics*, Brinkman F and Parkhill J (Eds.), 11:442–446. <http://doi:10.1016/j.mib.2008.09.011>

- van Berkum, P., & Fuhrmann, J.J. (2000). Evolutionary relationships among the soybean bradyrhizobia reconstructed from 16S rRNA gene and internally transcribed spacer region sequence divergence. *Int. J. Syst. Evol. Microbiol.*, 50, 2165-2172.
- van Berkum, P., & Fuhrmann, J.J. (2009). Evidence from Internally Transcribed Spacer Sequence Analysis of Soybean Strains that Extant *Bradyrhizobium* spp. Are Likely the Products of Reticulate Evolutionary Events. *Appl. Environ. Microbiol.*, 75 (1), 78-82. doi:10.1128/AEM.01408-08
- Vincent, J.M. (1970): A Manual for the Practical Study of the Root-Nodule Bacteria. Blackwell Scientific, Oxford
- Vinuesa, P., Rojas-Jiménez, K., Contreras-Moreira, B., Mahna, S.K., Prasad, B.N., Moe, H., Selvaraju, S.B., Thierfelder, H., & Werner, D. (2008). Multilocus Sequence Analysis for Assessment of the Biogeography and Evolutionary Genetics of Four *Bradyrhizobium* Species That Nodulate Soybeans on the Asiatic Continent. *Appl. Environ. Microbiol.*, 74 (22), 6987-6996. <http://doi:10.1128/AEM.00875-08>
- Vos, M., Quince, C., Pijl, A.S., de Hollander, M., & Kowalchuk, G.A. (2012). A Comparison of *rpoB* and 16S rRNA as Markers in Pyrosequencing Studies of Bacterial Diversity. *PLoS ONE* 7(2), 1-8, e30600. doi:10.1371/journal.pone.0030600
- Weatherbase. Retrieved from <http://www.weatherbase.com/weather/>
- Weaver, R.W., & Frederick LR. (1974). Effect of Inoculum Rate on Competitive Nodulation of *Glycine max* L. Merrill. II. Field Studies. *Agron. J.*, 66 (2), 233-236. doi:10.2134/agronj1974.00021962006600020015x

- Weisburg, W.G., Barm, S.M., Pelletier, D.A., & Lane, D.J. (1991). 16S ribosomal DNA amplification for phylogenetic study. *J. Bacteriol.*, 173, 697–703.
- Williams, L.F. & Lynch, D.L. (1954). Inheritance of a non-nodulation character in the soybean. *Agron. J.*, 46, 28–29.
- Wongphatcharachai, M., Staley, C., Wang, P., Moncada, K.M., Sheaffer, C.C., & Sadowsky, M.J. (2015a). Predominant populations of indigenous soybean-nodulating *Bradyrhizobium japonicum* strains obtained from organic farming systems in Minnesota. *J. Appl. Microbiol.*, 118 (5), 1152–1164. doi: 10.1111/jam.12771
- Wongphatcharachai, M., Wang, P., Staley, C.L., Ferguson, J.A., Moncada, K.M., Sheaffer, C.C., & Sadowsky, M.J. (2015b). Site-specific distribution and competitive ability of indigenous bean-nodulating rhizobia isolated from organic fields in Minnesota. *J. Biotechnol.*, 214, 158–168. doi: 10.1016/j.jbiotec.2015.09.009
- Yamakawa, T., Hussain, A.K.M.A., & Ishizuka J. (2003). Soybean preference for *Bradyrhizobium japonicum* for nodulation. *Soil Sci. Plant Nutr.*, 49 (6), 835–841. [http://doi: 10.1080/00380768.2003.10410345](http://doi:10.1080/00380768.2003.10410345)
- Yamakawa, T., & Saeki, Y. (2013). Inoculation Methods of *Bradyrhizobium japonicum* on Soybean in South-West Area of Japan, A Comprehensive Survey of International Soybean Research - Genetics, Physiology, Agronomy and Nitrogen Relationships, Prof. James Board (Ed.), ISBN: 978-953-51-0876-4, *InTech*, <http://doi:10.5772/52183>
- Yan, J., Han, X.Z., Ji, Z.J., Li, Y., Wang, E.T., Xie, Z.H., & Chen, W.F. (2014). Abundance and Diversity of Soybean-Nodulating Rhizobia in Black Soil Area

Impacted by Land Use and Crop Management. *Appl. Environ. Microbiol.*, 80 (17), 5394-5402. doi:10.1128/AEM.01135-14

Young, J.M. (2003). The genus name *Ensifer* Casida 1982 takes priority over *Sinorhizobium* Chen *et al.* 1988, and *Sinorhizobium morelense* Wang *et al.* 2002 is a later synonym of *Ensifer adhaerens* Casida 1982. Is the combination '*Sinorhizobium adhaerens*' (Casida 1982) Willems *et al.* 2003 legitimate? Request for an opinion. *Int. J. Syst. Evol. Microbiol.*, 53, 2107–2110. doi:10.1099/ij.s.0.02665-0

Zhang, Y.M., Li Y. Jr., Chen, W.F., Wang, E.T., Tian, C.F., Li, Q.Q., Zhang, Y.Z., Sui, X.H., & Chen, W.X. (2011). Biodiversity and Biogeography of Rhizobia Associated with Soybean Plants Grown in the North China Plain. *Appl. Environ. Microbiol.*, 77 (18), 6331–6342. doi: 10.1111/j.1574-6968.2008.01169.x

Zhao, L., Fan, M., Zhang, D., Yang, R., Zhang, F., Xu, L., Wei, X., Shen, Y., & Wei, G. (2014). Distribution and diversity of rhizobia associated with wild soybean (*Glycine soja* Sieb. & Zucc.) in Northwest China. *Syst. Appl. Microbiol.*, 37, 449–456. <http://dx.doi.org/10.1016/j.syapm.2014.05.011>

松浦 祥太. (2014). 土着ダイズ根粒菌の地理的分布と群集構造の比較解析. 宮崎大学大学院農学工学総合研究科資源環境科学専攻. 博士学位論文

城 惣吉. (2014). ダイズ根粒菌の環境適応機構と群集構造に関する研究. 宮崎大学大学院農学工学総合研究科資源環境科学専攻. 博士学位論文